

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 March 2007 (08.03.2007)

PCT

(10) International Publication Number
WO 2007/028118 A2

(51) International Patent Classification:
C07D 401/14 (2006.01) **C07D 417/14** (2006.01)

(21) International Application Number:
PCT/US2006/034406

(22) International Filing Date:
1 September 2006 (01.09.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/714,074 2 September 2005 (02.09.2005) US

(71) Applicant (*for all designated States except US*): **VISEN MEDICAL, INC.** [US/US]; 12B Cabot Road, Woburn, MA 01801 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **RAJOPADHYE, Milind** [IN/US]; 21 Honeysuckle Road, Westford, MA 01886 (US). **NARAYANAN, Narasimhachari** [US/US]; 73 Russells Way, Westford, MA 01886 (US). **PETERSON, Jeffrey, D.** [US/US]; 30 Hillando Drive, Shrewsbury, MA 01545 (US).

(74) Agents: **BROOK, David, E.** et al.; **HAMILTON, BROOK, SMITH & REYNOLDS, P.C.**, 530 Virginia Road, P.O. Box 9133, Concord, MA 01742-9133 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NICOTINIC ACID AND PICOLINIC ACID DERIVED NEAR-INFRARED FLUOROPHORES

(57) Abstract: This invention relates to new fluorescent chemical entities, including fluorescent molecules that comprise a carboxyl or carbonyl functionalized pyridine moiety. This invention also relates to the corresponding reactive versions of such molecules. This invention also relates to the corresponding conjugates with moieties such as peptides, proteins, various biomolecules, carbocyclic and heterocyclic compounds, sugars, and their uses thereof.



WO 2007/028118 A2

NICOTINIC ACID AND PICOLINIC ACID DERIVED NEAR-INFRARED FLUOROPHORES

RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No.
5 60/714,074, filed on September 2, 2005.

The entire teachings of the above application are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Optical imaging is an evolving clinical imaging modality that uses
10 penetrating light rays to create images. Light in the red and near-infrared (NIR)
range (600-1200 nm) is used to maximize tissue penetration and minimize
absorption from natural biological absorbers such as hemoglobin and water. (Wyatt,
Phil. Trans. R. Soc. London B **352**:701-706, 1997; Tromberg, et al., *Phil. Trans. R.*
Soc. London B **352**:661-667, 1997).

15 Besides being non-invasive, optical imaging methods offer a number of
advantages over other imaging methods: they provide generally high sensitivity, do
not require exposure of test subjects or lab personnel to ionizing radiation, can allow
for simultaneous use of multiple, distinguishable probes (important in molecular
imaging), and offer high temporal and spatial resolution (important in functional
20 imaging and in vivo microscopy, respectively).

In fluorescence imaging, filtered light or a laser with a defined bandwidth is
used as a source of excitation light. The excitation light travels through body
tissues. When it encounters a reporter molecule (i.e., contrast agent or imaging

has detectably different properties from the excitation light. The resulting emitted light then can be used to construct an image.

Most optical imaging techniques have relied on the use of organic and inorganic fluorescent molecules as the reporter molecule.

5 Fluorescent dyes are generally known and used for fluorescence labeling and detection of various biological and non-biological materials by procedures such as fluorescence microscopy, fluorescence immunoassay and flow cytometry. A typical method for labeling such materials with fluorescent dyes is to create a fluorescent complex by means of bonding between suitable groups on the dye molecule and
10 compatible groups on the material to be labeled. In this way, materials such as cells, tissues, amino acids, proteins, antibodies, drugs, hormones, nucleotides, nucleic acids, lipids and polysaccharides and the like may be chemically labeled and detected or quantified, or may be used as fluorescent probes which can bind specifically to target materials and detected by fluorescence detection methods.
15 Brightly fluorescent dyes permit detection or location of the attached materials with great sensitivity.


 Certain carbocyanine or polymethine dyes have demonstrated utility as labeling reagents for a variety of biological applications, e.g. U.S. Pat. No. 4,981,977 to Southwick, et al. (1991); 5,268,486 to Waggoner et al. (1993); U.S. Pat. No. 5,569,587 to Waggoner (1996); 5,569,766 to Waggoner et al. (1996); U.S. Pat. No. 5,486,616 to Waggoner et al. (1996); U.S. Pat. No. 5,627,027 to Waggoner (1997); U.S. Pat. No. 5,808,044 to Brush, et al. (1998); U.S. Pat. No. 5,877,310 to Reddington, et al. (1999); U.S. Pat. No. 6,002,003 to Shen, et al. (1999); U.S. Pat. No. 6,004,536 to Leung et al. (1999); U.S. Pat. No. 6,008,373 to Waggoner, et al.
20 (1999); U.S. Pat. No. 6,043,025 to Minden, et al. (2000); U.S. Pat. No. 6,127,134 to Minden, et al. (2000); U.S. Pat. No. 6,130,094 to Waggoner, et al. (2000); U.S. Pat. No. 6,133,445 to Waggoner, et al. (2000); also WO 97/40104, WO 99/51702, WO 01/21624, and EP 1 065 250 A1; and Tetrahedron Letters 41, 9185-88 (2000); all of the above incorporated by reference.

30 Comprehensive reviews regarding polymethine dyes have been by written by L. G. S. Brooker, "The Theory of the Photographic Process" Mees Ed., Macmillan, New York, (1942), p. 987 and (1966), p. 198; Frances M. Hamer, in "The Chemistry

of Heterocyclic Compounds", Vol 18, "The Cyanine Dyes and Related Compounds", Weissberger, Ed, Wiley Interscience, New York, (1964); G. E. Ficken, "The Chemistry of Synthetic Dyes", Vol 4, K. Venkataraman Ed., Academic Press, New York, (1971), p.211; A. I. Kiprianov, Usp. Khim., 29, 1336, (1960), 35, 361 (1966), 40, 594 (1971); D. W. Heseltine, "The Theory of the Photographic Process", 4th edition, James Ed., Macmillan, New York, (1977), chapter 8, "Sensitising and Desensitising Dyes"; S. Daehne, Phot. Sci. Eng., 12, 219 (1979); D. J. Fry, "Rodd's Chemistry of Carbon Compounds", "Cyanine Dyes and Related Compounds", Vol. IVb, chapter 15, p.369 Elsevier, Amsterdam, (1977); Supplement to Vol. IVb, 2nd Edition (1985), p.267; H. Zollinger, "Color Chemistry", VCH, Weinheim (1987), chapters 3 and 14; D. M. Sturmer, "The Chemistry of Heterocyclic Compounds", "Special Topics in Heterocyclic Chemistry", chapter VIII, "Synthesis and Properties of Cyanine and Related Dyes", Weissberger Ed., Wiley, New York, (1977); "The Kirk-Othmer Encyclopaedia of Chemical Technology" Vol 7, p. 782, "Cyanine Dyes", Wiley, New-York, (1993).

To be useful as a label, a dye has to be provided with a suitable side chain containing a functional group. The method and site of introduction of a side chain containing a functional group into the structure for the purpose of conjugation, or binding to another molecule, represents the innovative step in the inventions concerning the use of the dye as a labeling reagent. Typically, only one such functionalized side arm is used in order to avoid cross-linking or purification problems. One aspect in the design of polymethine labeling reagents has been to attach the functionalized side arm to one of the heterocyclic nuclei of the dye, formula (1):

25



See, for instance: J. S. Lindsey, P. A. Brown, and D. A. Siesel, "Visible Light-Harvesting in Covalently-Linked Porphyrin-Cyanine Dyes, Tetrahedron, 45, 4845, (1989); R. B. Mujumdar, L. A. Ernst, S. R. Mujumdar, and A. S. Waggoner, "Cyanine Dye Labelling Reagents Containing Isothiocyanate Groups", Cytometry, 10, 11 (1989); L. A. Ernst, R. K. Gupta, R. B. Mujumdar, and A. S. Waggoner,

30

- "Cyanine Dye Labelling Reagents for sulphhydryl Groups", Cytometry, 10, 3, (1989); P. L. Southwick P. L., L. A. Ernst, E. W. Tauriello, S. R. Parker, R. B. Mujumdar, S. R. Mujumdar, H. A. Clever, and A. S. Waggoner, "Cyanine Dye Labelling Reagents-Carboxymethylindocyanine Succinimidyl Esters", Cytometry 11, 418
- 5 (1990); R. B. Mujumdar, L. A. Ernst, Swati R. Mujumdar, C. J. Lewis, and A. S. Waggoner, "Cyanine Dye Labelling Reagents: Sulfoindocyanine Succinimidyl Esters", Bioconjugate Chemistry, 4, 105, (1993); A. J. G. Mank, E. J. Molenaar, H. Lingeman, C. Goojer, U. A. Th. Brinkman, and N. H. Velthorst, "Visible Diode Laser Induced Fluorescence Detection in Liquid Chromatography after Precolumn
- 10 Derivatisation of Thiols", Anal. Chem., 65, 2197, (1993); H. Yu., J. Chao, D. Patek, S. R. Mujumdar, and A. S. Waggoner, "Cyanine dye dUTP analogs for enzymatic labelling of DNA Probes", Nucl. Acids Res 22, 3226, (1994); Z. Zho, J. Chao, H. Yu, and A. S. Waggoner, "Directly labelled DNA probes using fluorescent nucleotides with different length linkers", Nucl. Acids, Res, 22, 3226. A. J. G.
- 15 Mank, H. T. C. van der Laan, , H. Lingeman, Cees Goojer, U. A. Th. Brinkman, and N. H. Velthorst, "Visible Diode Laser-Induced Fluorescence Detection in Liquid Chromatography after Precolumn Derivatisation of Amines", Anal. Chem., 67, 1742, (1995); S. R. Mujumdar, R. B. Mujumdar, C. M. Grant, and A. S. Waggoner, "Cyanine Labelling Reagents: sulfobenzoinocyanine succinimidyl esters",
- 20 Bioconjugate Chemistry, 7, 356, (1996). Patent Literature: P. L. Southwick, and A. S. Waggoner, "Intermediate for and Fluorescent Cyanine Dyes containing Carboxylic Acid Groups", U.S. Pat. No. 4,981,977, Jan. 1, 1991; A. S. Waggoner, L. A. Ernst, and Mujumdar, R. B., "Method for Labelling and Detecting Materials Employing Arylsulfonate Cyanine Dyes", U.S. Pat. No. 5,268,486, Dec. 7., 1993; A.
- 25 S. Waggoner, "Cyanine Dyes as Labelling Reagents for Detection of Biological and Other Materials by Luminescence Methods", U.S. Pat. No. 5,627,027, May 6, 1996; A. S. Waggoner, and R. B. Mujumdar, "Rigidised Trimethine Cyanine Dyes", WO99/311181; G.-Y. Shen, T. S. Dobashi, "Cyanine Dye Activating Group with Improved Coupling Selectivity"; T. S. G. M. Little, R. Raghavachari; N. Narayanan;
- 30 H. L. Osterman, "Fluorescent Cyanine Dyes", U.S. Pat. No. 6,027,709, Feb. 22, 2000.

The general synthetic strategy necessary to prepare these labeling reagents is as follows. First, a quaternized nitrogen heterocycle Z^1 is prepared. Then, this heterocyclic base is reacted with a polymethine linker (PML) that is an electrophilic reagent such as $\text{PhNH}-(\text{CH}=\text{CH})_n-\text{CH}=\text{NPh.HCl}$ or $\text{RO}-(\text{CH}=\text{CH})_n-\text{CH}(\text{OR})_2$,
5 where Ph is a phenyl ring and R a methyl or ethyl group, to obtain a so-called hemicyanine dye, $Z^1-(\text{CH}=\text{CH})_n\text{NPh}$ or $Z^1-(\text{CH}=\text{CH})_n\text{NAcPh}$, where Ac is the acetyl radical, or $Z^1-(\text{CH}=\text{CH})_n-\text{OR}$. These intermediates are then reacted with a different quaternary nitrogen heterocycle, Z^2 . The functionalized side arm can be attached either to the first or to the second quaternized nitrogen heterocycle.
10 The hemicyanine intermediates, however, can be difficult to obtain in good yields and/or in a pure form (see, for example, F. M. Hamer, "Some Unsymmetrical Pentamethincyanine Dyes and their Tetramethin Intermediates", J. Chem. Soc., 32 (1949) and R. B. Mujumdar, L. A. Ernst, Swati R. Mujumdar, C. J. Lewis, and A. S. Waggoner, "Cyanine Dye Labelling Reagents: Sulfoindocyanine Succinimidyl
15 Esters", Bioconjugate Chemistry, 4, 105, (1993).


Thus polymethine dyes that are efficient and easy to produce as well as suitable for preparing conjugates with biomolecules are desirable.

SUMMARY OF THE INVENTION

20 Fluorescent dye compounds with a polymethine linker bridge that are bright, highly fluorescent dyes that absorb and emit between about 440 and about 1100 nm, have now been discovered.

The present invention is directed to fluorescent dye compounds with a polymethine linker bridge that has been modified by a carboxyl containing pyridine
25 ring which are capable of producing strong fluorescence in the 600 nm to 900 nm region of the spectrum and also contain functional groups and/or solubilizing groups which render the dye suitable for covalent labeling, in particular to biological molecules and other target materials.

In one embodiment, the present invention is directed to a polymethine
30 fluorochrome compound represented by formulae (2) or a salt thereof:



(2)

X is independently selected from the group consisting of C(CH₂Y₁)(CH₂Y₂), O, S, and Se;

5 Y₁ and Y₂ are independently selected from the group consisting of H, C₁-C₂₀ aliphatic group and a C₁-C₂₀ aliphatic group substituted with -OR*, N(R*)₂ or -SR*;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;

R₁ is selected from the group consisting of H, (CH₂)_xCH₃, (CH₂)_nSO₃⁻ and
 10 (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R₄ is selected from the group consisting of H, (CH₂)_xCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

15 R₂ and R₃ are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group.

20 In another embodiment present invention is directed to a polymethine fluorochrome compound represented by formulae (2) or a salt thereof when:

X is independently selected from the group consisting of C(CH₂Y₁)(CH₂Y₂), O, S, and Se;

Y₁ and Y₂ are independently selected from the group consisting of H, C₁-C₂₀ aliphatic group and a C₁-C₂₀ aliphatic group substituted with -OR*, N(R*)₂ or -SR*;;
 25

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;

R_1 is selected from the group consisting of $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_4 is selected from the group consisting of $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_2 and R_3 are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group.

In another embodiment, the present invention is directed to a biocompatible fluorescent molecule comprising one or more biomolecules chemically linked to a compound of the present invention.

In another embodiment, the present invention is directed to a biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound of the present invention, wherein the compound has an absorption and emission maxima between about 500 nm and about 900 nm.

In another embodiment, the present invention is directed to a biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound of the present invention wherein the biocompatible fluorescent molecule is activated after target interaction.

In another embodiment, the present invention is directed to a biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound of the present invention wherein the biocompatible fluorescent molecule has a high binding affinity to a target.

In another embodiment, the present invention is directed to a method of in vivo optical imaging, the method comprising:

(a) administering to a subject a compound or biocompatible fluorescent molecule of the present invention

- (b) allowing time for the compound or biocompatible fluorescent molecule to distribute within the subject or to contact or interact with a biological target;
- (c) illuminating the subject with light of a wavelength absorbable by the compound or biocompatible fluorescent molecule; and
- (d) detecting the optical signal emitted by the compound or biocompatible fluorescent molecule.

In another embodiment, the present invention is directed to a biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound of the present invention, wherein the biocompatible fluorescent molecule is a labeled cell.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a fluorescence image of a compound of the present invention in tumors of a female NU/NU mice (6-8 weeks old) after 24 hrs on a fluorescence reflectance system (FRI, Kodak 2000MM) system.

DETAILED DESCRIPTION OF THE INVENTION

A description of preferred embodiments of the invention follows.

The present invention is directed to bright, highly fluorescent compounds (dyes) that absorb and/or emit between about 440 and about 1100 nm, between about 550 and about 800nm, between about 500 and about 900 nm or between about 600 and about 900 nm and conjugates thereof. It will be appreciated that compounds (fluorochromes) with excitation and emission wavelengths in other spectrums, such as the visible and ultraviolet light spectrum, are also encompassed by the present invention.

The structures of the compounds of the present invention, in general are based on nicotinic acid and picolinic acid derivatives that confer high quantum yields of fluorescence. Moreover, in certain embodiments of the present invention the compounds contain functional or reactive groups which may be used to chemically link with complementary groups on target molecules.

“Chemically linked” or “chemically link” means connected by an attractive force between atoms strong enough to allow the combined aggregate to function as a unit. This includes, but is not limited to, chemical bonds such as covalent bonds, non-covalent bonds such as ionic bonds, metallic bonds, and bridge bonds, hydrophobic interactions, hydrogen bonds, and van der Waals interactions. This also includes crosslinking or caging.

The term “compounds,” as used herein, refers to “polymethine fluorochromes”, “fluorochromes”, “fluorescent dyes”, “cyanine dyes” “carbocyanine dyes” and “dyes” of the instant invention. These terms are used interchangeably to refer to the compounds of the instant invention.

In one embodiment, the compounds of the invention comprise two heterocyclic ring systems bound together by a polymethine linker (PML), according to the formula (1):




wherein Z^1 is a heterocyclic ring system such as an indolinium ring, Z^2 is a second heterocyclic ring system, such as, an indolinium ring, and PML is a polymethine linker that is substituted with a carboxyl containing heterocyclic ring, such as, pyridine. The Z^1 and Z^2 ring systems are optionally further substituted by a variety of substituents or are fused to additional rings that are optionally further substituted.

In one aspect, the compounds of the present invention are further substituted one or more times by sulfo or sulfoalkyl. By "sulfo" is meant sulfonic acid, or salt of sulfonic acid (sulfonate). Similarly, by "carboxyl" is meant carboxylic acid, carboxylate ester or salt of carboxylic acid. "Phosphate" is an ester of phosphoric acid, and includes salts of phosphate. "Phosphonate" means phosphonic acid and includes salts of phosphonate. Similarly for “carbonyl” groups such as, but not limited to carbonyl halide, (e.g., chloride) and carboxamide are included. As used herein, unless otherwise specified, the alkyl portions of substituents such as alkyl, alkoxy, arylalkyl, alkylamino, dialkylamino, trialkylammonium, or perfluoroalkyl are optionally saturated, unsaturated, linear or branched, and all alkyl, alkoxy,


alkylamino, and dialkylamino substituents are themselves optionally further substituted by carboxy, sulfo, amino, or hydroxy.

In one embodiment the present invention is directed to a compound represented by formulae (2) or a salt thereof, wherein:

5




- wherein X is independently selected from C(CH₂Y₁)(CH₂Y₂), O, S, Se; and Y₁ and Y₂ are independently selected from H and C₁-C₂₀ alkyl that is linear or branched, saturated or unsaturated, optionally containing N, S, O in various forms; W represents non-metal atoms required to form a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring. R₁ is selected from the group consisting of H, (CH₂)_nCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein n is an integer selected from 0 to 6 when R₁ is (CH₂)_nCH₃, and n is an integer selected from 2 to 6 when R₁ is (CH₂)_nSO₃⁻ or (CH₂)_nSO₃H. R₄ is selected from the group consisting of H, (CH₂)_nCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein n is an integer selected from 0 to 6 when R₄ is (CH₂)_nCH₃, and n is an integer selected from 2 to 6 when R₄ is (CH₂)_nSO₃⁻ or (CH₂)_nSO₃H. R₂ and R₃ are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety. Q is selected from a group consisting of a carboxyl functionalized nitrogen containing heterocyclic ring. Q is selected from a group consisting of a carbonyl functionalized nitrogen containing 6-membered heterocyclic ring, such as pyridine. Q is selected from a group consisting of isonicotinic acid, nicotinic acid and picolinic acid. Q is selected from the groups shown:



Herein, the carboxyl group is also in the form of an ester, an activated ester or carbonyl halide that is capable of reacting with nucleophiles. The carboxyl group is also in a form selected from the group CO-Obenzotriazolyl, CO-O-N-succinimidyl, CO-Otetrafluorophenyl, CO-Opentafluorophenyl, CO-Oimidazolyl, CO-Op-nitrophenyl.


Such molecules of formula (2) are chemically linked to biocompatible fluorescent molecules for a variety of applications including in vivo imaging.

In another embodiment the present invention is directed to a compound represented by formulae (2) or a salt thereof, wherein:



wherein X is independently selected from C(CH₂Y₁)(CH₂Y₂), O, S, Se; and Y₁ and Y₂ are independently selected from H and C₁-C₂₀ alkyl that is linear or branched, saturated or unsaturated, optionally containing N, S, O in various forms; W represents non-metal atoms required to form a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring. R₁ is selected from the group consisting of H, (CH₂)_nCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein n is an integer selected from 0 to 6 when R₁ is (CH₂)_nCH₃, and n is an integer selected from 2 to 6 when R₁ is (CH₂)_nSO₃⁻ or (CH₂)_nSO₃H. R₄ is selected from the group consisting of (CH₂)_nCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein n is an integer selected from 0 to 6 when R₄ is (CH₂)_nCH₃, and n is an integer selected from 2 to 6 when R₄ is (CH₂)_nSO₃⁻ or (CH₂)_nSO₃H. R₂ and R₃ are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety. Q is selected from a

group consisting of a carboxyl functionalized nitrogen containing heterocyclic ring. Q is selected from a group consisting of a carbonyl functionalized nitrogen containing 6-membered heterocyclic ring, such as pyridine. Q is selected from a group consisting of isonicotinic acid, nicotinic acid and picolinic acid. Q is selected from the groups shown:



Herein, the carboxyl group is also in the form of an ester, an activated ester or carbonyl halide that is capable of reacting with nucleophiles. The carboxyl group is also in a form selected from the group CO-Obenzotriazolyl, CO-O-N-succinimidyl, CO-Otetrafluorophenyl, CO-Opentafluorophenyl, CO-Oimidazolyl, CO-Op-nitrophenyl.

Such molecules of formula (2) are chemically linked to biocompatible fluorescent molecules for a variety of applications including in vivo imaging.

In one embodiment for structural formula (2) Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl or 6-membered heteroaryl ring substituted with a carbonyl group.

In certain embodiments:

- a) the carboxyl substituent on Q is selected from the group consisting of an ester, an activated ester; or
- b) the carboxyl substituent on Q is selected from the group consisting of CO-Obenzotriazolyl, CO-ON-succinimidyl, CO-Otetrafluorophenyl, CO-Opentafluorophenyl, CO-Oimidazolyl, and CO-Op-nitrophenyl; and
- c) the carbonyl substituent on Q is in the form of a carbonyl halide.

In another embodiments, Q is a carboxyl substituted nitrogen containing heterocyclic ring.


In another embodiments, Q is selected from the group consisting of carboxyl substituted pyridine, pyrimidone, pyrazine, and pyridazine.

In another embodiments, Q is carboxyl substituted pyridine.

In another embodiments, Q is selected from a group consisting of isonicotinic acid, nicotinic acid and picolinic acid.

In another embodiments, Q is represented by a structural formula selected from a group consisting of:

5



In another embodiments, Q is a carbonyl substituted nitrogen containing 6-membered heteroaryl ring


In another embodiments, Q is a carbonyl substituted pyridine.

10

In the immediately preceding nine embodiments, the terms heterocyclic ring and heteroaryl ring both refer to heteroaryl ring as defined herein, such as pyridine.

In another embodiment, the present invention provides compounds representing Z^1 -(PML)- Z^2 by the formula (2):

15



wherein, W represents non-metal atoms required to form a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring.

20


In one aspect W represents the atoms necessary to form one to two fused aromatic rings having 6 atoms in each ring, which atoms are selected from -CH, -C, -CR₇, and -NR₈, where R₈ is 0 or 1 (such that each ring nitrogen is either quaternized or not), and each R₇ independently contains sulfo, trifluoromethyl, or halogen; R₈ independently contains a C₁-C₈ alkyl, in turn containing independently an H, amino or sulfo.

25

Incorporation of one or more non-hydrogen substituents on the fused rings can be used to tune the absorption and emission spectrum of the resulting dye.

Selected examples of the basic structure of Z^1 and Z^2 are shown below. These basic structures (3- 6) are optionally further substituted as described in this section.

5



In one embodiment, X is independently selected from O, S, Se, -
 10 $C(CH_2Y_1)(CH_2Y_2)$, wherein Y_1 and Y_2 are independently selected from H and C_1 -
 C_{20} alkyl that is linear or branched, saturated or unsaturated, optionally containing
 N, S, O in various forms,

In another embodiment Y_1 and Y_2 together are part of a cyclic ring; or X is
 independently selected from $-CR_5R_6$, wherein R_5 and R_6 , which may be the same or
 15 different, are alkyl, cycloalkyl, or arylalkyl, together part of a cyclic system and
 optionally further substituted.

In one aspect of the invention, R_2 and R_3 taken in combination complete a
 five or six- membered ring.

The substituent R_1 is typically selected from the group consisting of
 20 $(CH_2)_nCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein n is an integer selected from 0 to
 6 when R_1 is $(CH_2)_nCH_3$, and n is an integer selected from 2 to 6 when R_1 is
 $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$. In one aspect, R_1 is $(CH_2)_3SO_3^-$ or $(CH_2)_4SO_3^-$ or
 CH_2CH_3 . In one aspect of the invention R_1 is a substituent containing an aryl
 sulfonate or an amino group or a phthalimido group.

25 The substituent R_1 is typically selected from the group consisting of H,
 $(CH_2)_nCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein n is an integer selected from 0 to
 6 when R_1 is $(CH_2)_nCH_3$, and n is an integer selected from 2 to 6 when R_1 is
 $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$. In one aspect, R_1 is $(CH_2)_3SO_3^-$ or $(CH_2)_4SO_3^-$ or
 CH_2CH_3 . In one aspect of the invention R_1 is a substituent containing an aryl
 30 sulfonate or an amino group or a phthalimido group.

The substituent R_4 is typically selected from the group consisting of $(CH_2)_nCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein n is an integer selected from 0 to 6 when R_4 is $(CH_2)_nCH_3$, and n is an integer selected from 2 to 6 when R_4 is $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$. In one aspect, R_4 is $(CH_2)_3SO_3^-$ or $(CH_2)_4SO_3^-$ or CH_2CH_3 . In one aspect of the invention R_4 is a substituent containing an aryl sulfonate or an amino group or a phthalimido group.


The substituent R_4 is typically selected from the group consisting of H, $(CH_2)_nCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein n is an integer selected from 0 to 6 when R_4 is $(CH_2)_nCH_3$, and n is an integer selected from 2 to 6 when R_4 is $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$. In one aspect, R_4 is $(CH_2)_3SO_3^-$ or $(CH_2)_4SO_3^-$ or CH_2CH_3 . In one aspect of the invention R_4 is a substituent containing an aryl sulfonate or an amino group or a phthalimido group.

The substituents R_2 and R_3 are independently selected from the group H, halogens, carboxylate, carboxylic acid, carboxylic esters, amino, amide, alkyl or aryl sulfonamide, hydroxy, alkoxy, aryloxy, sulfate, cyano, nitro, azido, alkylamino, dialkylamino, trialkylammonium, phosphate, phosphate ester, phosphonate, sulphonic acid and a sulphonate moiety. In certain embodiments, R_2 and R_3 are independently, sulphonic acid or a salt thereof.

In one aspect of the invention, R_2 and R_3 could imply per-substitution, as in per-fluorinated W. Per-fluorination or poly-fluorination can lead to enhancement of fluorescence quantum yield.

As used herein "enhancement" means an increase in the fluorescence quantum yield, by about 5%, about 10%, about 15%, about 25%, about 50%, about 80%, about 90% about 95% about 98% about 99% about 100%.

In one aspect of the invention, the formula $Z^1-(PML)-Z^2$ is represented according to the formula (7):



In one aspect of the invention, the compounds of the invention are sulfonated one or more times. If the compound of the invention is substituted by sulfo, it is

typically sulfonated at R_2 or R_3 or both, (that is, for example, R_2 and/or R_3 are a sulfonic acid moiety, sulfonate moiety or sulfonamamide) or sulfoalkylated at R_1 or R_4 or both (that is, for example, R_1 and/or R_4 are $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$), or is both sulfonated and sulfoalkylated. In a particular embodiment, the compounds are
5 are sulfonated up to three times (at positions corresponding to R_2 and R_3 , and as sulfoalkyl at one of R_1 or R_4), leaving one of R_1 or R_4 for the location of a reactive group.


As used herein the terms sulfonic acid and $(CH_2)_nSO_3H$ or a sulfonate group and $(CH_2)_nSO_3^-$ can be used interchangeably. However in certain embodiments the
10 terms a sulfonic acid moiety, sulfonate moiety or sulfonamamide refer to substituents which are attached to the remainder of the molecule by the a sulfonic acid moiety, sulfonate moiety or sulfonamamide moiety, ie., $-SO_2NR'R''$

In certain embodiments, at least one of R_1 to R_3 is or contains a sulphonic acid moiety or a sulphonate moiety. In certain embodiments, at least one of R_1 to R_3
15 is a sulphonic acid moiety or a sulphonate moiety.

In certain embodiments, R_1 and R_4 are independently $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$.

In one embodiment of the present invention the compounds are sulfonated up to four times (at R_2 and R_3 , and as sulfoalkyl at R_1 and R_4). In another embodiment
20 of the present invention the compounds are sulfonated at least four times (at R_2 and R_3 , and as sulfoalkyl at R_1 and R_4). In another embodiment of the present invention, any reactive group (or chemically linked molecule) can be attached at Q, in addition the compounds of the present invention being sulfonated up to four times (at R_2 and R_3 , and as sulfoalkyl at R_1 and R_4). In another embodiment of the present invention,
25 any reactive group (or chemically linked molecule) can be attached at Q, in addition the compounds of the present invention being sulfonated at least four times (at R_2 and R_3 , and as sulfoalkyl at R_1 and R_4). This extra sulfonation, as well as the change in attachment site, results in reactive dyes and dye conjugates that are brighter, more soluble in aqueous solutions, and more resistant to the fluorescence quenching that
30 results from dye-dye stacking interactions.


In one aspect of the invention, the formula $Z^1-(PML)-Z^2$ is represented according to the formula (8):



In one aspect of the invention, the compounds of the invention that are
 5 sulfonated one or more times are typically sulfonated at R₂ or R₃, or sulfoalkylated
 at R₁ or R₄ or both, or is both sulfonated and sulfoalkylated. In particular, the
 compounds of the present invention are sulfonated up to four times leaving one of
 R₁ or R₄ for the location of a reactive group.

In one embodiment of the present invention the compounds are sulfonated up
 10 to six times (at R₂ and R₃, and as sulfoalkyl at R₁ and R₄). In another embodiment
 of the present invention the compounds are sulfonated at least six times (at R₂ and
 R₃, and as sulfoalkyl at R₁ and R₄). In another embodiment of the present invention,
 any reactive group (or chemically linked molecule) can be attached at Q, in addition
 the compounds of the present invention being sulfonated up to six times (at R₂ and
 15 R₃, and as sulfoalkyl at R₁ and R₄). In another embodiment of the present invention,
 any reactive group (or chemically linked molecule) can be attached at Q, in addition
 the compounds of the present invention being sulfonated at least six times (at R₂ and
 R₃, and as sulfoalkyl at R₁ and R₄). This extra sulfonation, as well as the change in
 attachment site, results in reactive dyes and dye conjugates that are brighter, more
 20 soluble in aqueous solutions, and more resistant to the fluorescence quenching that
 results from dye-dye stacking interactions.

In one embodiment, the PML moiety has the formula (9):



25

wherein Q is selected from a group consisting of carboxyl functionalized
 heterocyclic ring (heteroaryl substituted with a carboxy group). In this embodiment,

heterocyclic ring and heteroaryl ring refers to heteroaryl ring as defined herein, such as pyridine and functionalized refers to substituted.

Suitable examples of appropriate PML moieties for compounds of the present invention (carbocyanine dyes) have been described in the literature, including PML moieties that incorporate nonhydrogen substituents, ring structures, and rigidizing elements (U.S. Pat. No. 5,831,098 to Ollmann, Jr (1998); U.S. Pat. No. 6,086,737 to Patonay et al. (2000); U.S. Pat. No. 6,048,982 to Waggoner (2000); and U.S. Pat. No. 5,453,505 to Lee et al. (1995); U.S. Pat. No. 5,639,874 to Middendorf et al. (1997); U.S. Pat. No. 3,864,644 to Lincoln et al (1975); U.S. Pat. No. 4,011,086 to Simson (1977); U.S. Pat. No. 6,747,159 to Caputo (2004); all incorporated herein by reference in their entirety).

In one embodiment the invention is directed to a PML of the formula (9), wherein Q is selected from a group consisting of a functionalized nitrogen-containing heterocyclic ring. One aspect of the invention is a PML of the formula (9), wherein Q is selected from a group consisting of a substituted nitrogen-containing heteroaryl ring. In these embodiments, heterocyclic ring and heteroaryl ring refers to heteroaryl ring as defined herein, such as pyridine, and functionalized refers to substituted.

In one embodiment of the invention, Q contains at least one substituent which is a reactive group that is attached to the heterocyclic (heteroaryl) ring (Q) by a covalent linkage. In one embodiment the compounds of the present invention which contain a reactive group label a wide variety of organic or inorganic substances that contain or are modified to contain functional groups with suitable reactivity, resulting in chemical attachment of the compounds of the present invention to form a conjugated substance.

As used herein, "reactive group" means a moiety on a compound of the present invention or that can be added to a compound of the present invention that is capable of chemically reacting with a functional group on a different compound to form a covalent linkage, or a moiety on a different compound that is capable of chemically reacting with a functional group on compound of the present invention to form a covalent linkage. Typically the reactive group is an electrophile or nucleophile that can form a covalent linkage through exposure to the corresponding

functional group that is a nucleophile or electrophile, respectively. Alternatively, the reactive group is a photoactivatable group, and becomes chemically reactive only after illumination with light of an appropriate wavelength. Typically, the conjugation reaction between the compound of the present invention and the substance to be conjugated results in one or more atoms of the reactive group to be incorporated into a new linkage attaching the dye to the conjugated substance.


One aspect of the invention is a PML of the formula (9), wherein Q is selected from a group consisting of carboxyl functionalized nitrogen containing heterocyclic ring (heteroaryl substituted with carboxyl). In this embodiment, heterocyclic ring and heteroaryl ring refers to heteroaryl ring as defined herein, such as pyridine and functionalized refers to substituted.

In one aspect of the invention PML is of the formula (9), wherein Q is selected from a group consisting of carboxyl functionalized nitrogen containing 6-membered heterocyclic ring (heteroaryl substituted with carboxyl), such as pyridine, pyrimidone, pyrazine, and pyridazine. (heteroaryl substituted with carboxyl). In this embodiment, heterocyclic ring and heteroaryl ring refers to heteroaryl ring as defined herein, such as pyridine and functionalized refers to substituted.

Another aspect of the invention is a PML of the formula (9), wherein Q is selected from a group consisting of carboxyl functionalized nitrogen containing 6-membered heterocyclic ring, (heteroaryl substituted with carboxyl), such as pyridine. (heteroaryl substituted with carboxyl). In this embodiment, heterocyclic ring and heteroaryl ring refers to heteroaryl ring as defined herein, such as pyridine and functionalized refers to substituted.


In another aspect of the invention PML is of the formula (9), wherein Q is selected from a group consisting of nicotinic acid and picolinic acid or a salt thereof.

In one aspect of the invention is a PML of the formula (9), wherein Q is selected from the groups shown:




or a salt thereof.

- 5 In another aspect of the invention is a PML of the formula (9), wherein Q is selected from the groups shown:




- 10 wherein, the carboxyl group is also in the form of an activated ester (R₁₅) or carbonyl halide (R₁₆ = F, Cl, Br) that is capable of reacting with nucleophiles. The carboxyl group CO-OR₁₅ is also in a form selected from the group CO-Obenzotriazolyl, CO-ON-hydroxysuccinimidyl, CO-Otetrafluorophenyl, CO-Opentafluorophenyl, CO-Oimidazolyl, CO-Op-nitrophenyl.
- 15 The PML moiety typically originates from the coupling agent used in the synthesis of a compound of the present invention. For example, N,N'-diphenylformamidine and triethylorthoformate yields PML moieties. Malonaldehyde bis(phenylimine) hydrochloride, 1,1,3-trimethoxypropane, and 1,1,3,3-tetramethoxypropane and glutaconaldehyde dianil monochloride also yield PML
- 20 moieties (dyes).

In one aspect of the invention, the PML moiety is introduced into the dye using the malonodialdehydes moieties shown below:




Accordingly, the present invention provides compounds representing Z^1 -
(PML)- Z^2 by the formula (2):




wherein R_1 is $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$.

In one aspect the present invention, is directed to compounds having any of
10 the formulae (10-11):




(10)




(11)

In another aspect the present invention is directed to compounds having any
of the formulae (12-15):


20




(12)



(13)



(14)




(15)


5 In one aspect of the invention when Q is or contains an activated ester, the compound can be chemically linked to bifunctional linkers such as aminoethylmaleimide, aminopropylmaleimide, aminopropylazide, aminopropylthiol, mercaptoethylamine, propargylamine 3-aminopropanol, diaminopropane, and
 10 diaminobutane to provide additional reactive functional groups in a suitable solvent under neutral or basic conditions.

In one aspect of the invention when Q is or contains NH_2 , the compound of the invention can be chemically linked to bifunctional linkers such as propargylic acid, succinimidylpyridinedithiopropionate, maleimide-PEG-N-hydroxysuccinimide
 15 ester to provide additional reactive functional groups in a suitable solvent under neutral or basic conditions.

When a compound of the invention is depicted herein by structure indicating the positions of the double bonds in the rings an polymethine linker, it is to be understood that the structure also encompasses any resonance structures as shown, for example, in the figure below:



In one embodiment, the present invention provides compounds representing $\text{Z}^1\text{-(PML-BM)-Z}^2$ by the formula (16):



wherein BM a biomolecule chemically linked via Q residing on the PML.

In one aspect BM is a moiety that changes or alters or enhances accumulation, biodistribution, elimination, targeting, binding, and/or recognition of the compounds (fluorochromes) of the general structure Z^1 -(PML)- Z^2 . BMs include but are not limited to antibodies and fragments thereof, proteins, peptides, amino acids, antibodies (or antigen-binding antibody fragments, such as single chain antibodies), glycoproteins, ligands for cell receptors, polysaccharides, nucleosides, aptamers, cell receptors themselves, enzyme substrates, enzyme cofactors, biotin, hormones, neurohormones, neurotransmitters, growth factors, cytokines, lymphokines, lectins, selectins, toxins, and carbohydrates. Other targeting and delivery approaches using various biomolecules can also be used, such as folate-mediated targeting (Leamon & Low, *Drug Discovery Today*, 6:44-51, 2001), transferrin, vitamins, carbohydrates and ligands that target internalizing receptors, including, but not limited to, asialoglycoprotein receptor, somatostatin, nerve growth factor, oxytocin, bombesin, calcitonin, arginine vasopressin, angiotensin II, atrial natriuretic peptide, insulin, glucagons, prolactin, gonadotropin, various opioids and urokinase-type plasminogen activator. Also included are membrane, transmembrane, and nuclear translocation signal sequences, which can be derived from a number of sources including, without limitation, viruses and bacteria. BM can also be an organic molecule, polymer, dendrimer, drug, lipid, lipid assembly, therapeutic drug molecules, polymeric microparticle, cells, or a nanoparticle. In certain embodiments, BMs can also include small molecule drugs, phototherapeutic molecules and derivatives thereof.

In certain embodiments of the present invention, when BM is chemically linked to a compound of the present invention the fluorescence of the compound of the present invention is enhanced. In certain embodiments the fluorescence is

enhanced by about 10% , about 25%, about 50% or more than about 50% which compared with the unlinked compound.

In one aspect of the invention several copies of BM are chemically linked to Q via multivalent linkers or linkers containing several reactive functional groups to form a biocompatible fluorescent molecule of the structure $(Z^1-(PML)-Z^2)-((L)_w-(BM)q)_t$, wherein L is a linker or multivalent linker, and $t=1-6$, $w=1-500$ and $q=1-500$. $(L)_w$ represents copies of the same linker or a combination of different linkers. Suitable linkers include but are not limited to aminocaproic acid, glutamic acid, and poly-glutamic acid.

10 In another aspect of the invention, more than one compound (fluorochrome) of the structure $Z^1-(PML)-Z^2$ can be chemically linked to a single BM to form a biocompatible fluorescent molecule of the structure $[Z^1-(PML)-Z^2]_k-BM$, wherein $k=1-500$.

In one embodiment $Z^1-(PML-BM)-Z^2$ is a biocompatible fluorescent molecule wherein the compound has an absorption and emission maxima between about 440 and about 1100 nm, between about 550 and about 800nm, between about 500 and about 900 nm or between about 600 and about 900 nm.

In one embodiment the compounds (fluorochromes) and biocompatible fluorescent molecules of the present invention are activated after target interaction. "Activated after target interaction" is meant a change that alters a detectable property, *e.g.*, an optical property, of the fluorochromes or biocompatible fluorescent molecules. This includes, but is not limited to, a modification, alteration, or binding (covalent or non-covalent) that results in a detectable difference in properties, *e.g.*, optical properties of the fluorochromes or biocompatible fluorescent molecules, *e.g.*, changes in the fluorescence signal amplitude (*e.g.*, dequenching and quenching), change in wavelength, fluorescence lifetime, spectral properties, or polarity. In another embodiment, a quencher molecule is used to quench the fluorescent signal of the biocompatible fluorescent molecules. By adopting these activated and unactivated states, it is therefore possible to determine whether the fluorochrome or biocompatible fluorescent molecule is active or inactive in a subject by identifying a change in the signal intensity. In addition, the fluorochromes and biocompatible fluorescent molecules can be designed such that they exhibit little or no signal

until activated. Activation can be, without limitation, by enzymatic cleavage, enzymatic conversion, phosphorylation or dephosphorylation, conformation change due to binding, enzyme-mediated splicing, enzyme-mediated transfer, hybridization of complementary DNA or RNA, analyte binding, such as association with an
5 analyte such as Na^+ , K^+ , Ca^{2+} , Cl^- , or another analyte, change in hydrophobicity of the environment and chemical modification.

In one embodiment the compounds (fluorochromes) and biocompatible fluorescent molecules of the present invention have a high binding affinity to a target.

10 Also provided herein is a method of in vivo optical imaging, the method comprising (a) administering to a subject a compound (fluorochrome) or biocompatible fluorescent molecule of the present invention; (b) allowing time for the compound (fluorochrome) or biocompatible fluorescent molecule to distribute within the subject or to contact or interact with a biological target; (c) illuminating
15 the subject with light of a wavelength absorbable by the compound (fluorochrome) or biocompatible fluorescent molecule; and (d) detecting the optical signal emitted by the compound (fluorochrome) or biocompatible fluorescent molecule.

The optical signal generated by the compounds (fluorochromes) or biocompatible fluorescent molecules of the present invention, whether collected by
20 tomographic, reflectance, planar, endoscopic, microscopic, surgical goggles, video imaging technologies, or other methods such as microscopy including intravital and two-photon microscopy, and whether used quantitatively or qualitatively, is also considered to be an aspect of the invention.

One aspect of the invention is a method wherein the presence, absence,
25 distribution, or level of optical signal emitted by the compound (fluorochrome) or biocompatible fluorescent molecule is indicative of a disease state.

The invention also features a method of using the compounds (fluorochromes) and biocompatible fluorescent molecules of the present invention to detect an abnormality in a patient or subject, e.g., any abnormality associated with a
30 disease such as cancer, a cardiovascular disease, AIDS, a neurodegenerative disease, an inflammatory disease, a respiratory disease, a metabolic disease, a bone disease or an immunologic disease. The invention also features a method of assessing the

effect of a compound or therapy on a specified molecular target by using the compounds (compositions) of the present invention, wherein the subject is imaged prior to and after treatment with the compound or therapy, and the corresponding images are compared.

5 The imaging method steps of the present invention can also be repeated at predetermined intervals thereby allowing for the evaluation of emitted signal of the Z^1 -(PML)- Z^2 containing compound in a subject or sample over time. The emitted signal may take the form of an image. The subject may be a vertebrate animal, for example, a mammal, including a human. The animal may also be non-vertebrate,
10 (e.g., *C. elegans*, *drosophila*, or other model research organisms, etc.). The sample can include, without limitation, cells, cell culture, tissue sections, organs, organ sections, cytospin samples, or the like.

 The invention also features an in vivo method for selectively detecting and imaging two or more Z^1 -(PML)- Z^2 containing compounds simultaneously. The
15 method comprises administering to a subject two or more Z^1 -(PML)- Z^2 containing compounds, either at the same time or sequentially, whose optical properties are distinguishable. The method therefore allows the recording of multiple events or targets.

 The invention also features an in vivo method for selectively detecting and
20 imaging one or more Z^1 -(PML)- Z^2 containing compounds, simultaneously with one or more targeted or activatable optical imaging probes, or in a dual imaging protocol with magnetic resonance imaging, computed tomography (CT), X-ray, ultrasound, or nuclear medicine imaging modalities and their respective imaging agents. The method comprises administering to a subject one or more imaging probes, either at
25 the same time or sequentially, including at least one Z^1 -(PML)- Z^2 containing compound, whose properties are distinguishable from that of the others. In one aspect a dual imaging protocol is optical and magnetic resonance imaging using Z^1 -(PML)- Z^2 containing compounds sequentially or nearly simultaneously with magnetic resonance imaging agents, (for example, iron oxide based agents or
30 gadolinium based agents such as gadopentetate). The method therefore, allows the recording of multiple events or targets using more than one imaging modality or imaging agent.

In another aspect, the invention features an in vitro optical imaging method comprising contacting the sample with Z^1 -(PML)- Z^2 containing compounds; allowing time for the probes to become activated or bind to a target of interest in the sample; optionally, removing the unbound probes; illuminating the target with light
5 of a wavelength absorbable by the Z^1 -(PML)- Z^2 containing compounds; and detecting the optical signal emitted by the Z^1 -(PML)- Z^2 containing compounds.

After administration, detection can occur, for example, by in vitro methods, i.e., flow cytometry or by in vivo imaging methods, i.e., tomographic, catheter, planar/reflectance systems or endoscopic systems.

10 In one embodiment, the Z^1 -(PML)- Z^2 containing compounds derived thereof can be used to label a sample ex vivo. The sample, e.g., cells, can be derived directly from a subject or from another source (e.g., from another subject, cell culture etc.). The Z^1 -(PML)- Z^2 containing compound can be mixed with the cells to effectively label the cells (covalently or non-covalently) and the resulting labeled
15 cells injected into a subject. This method can be used for monitoring trafficking and localization of certain cell types, including T-cells, tumor cells, immune cells, stem cells, and other cell types. In particular, this method may be used to monitor cell based therapies. The sample can also be derived from non-mammalian sources including but not limited to plants, insects, viruses, bacteria, and bacteriophage.

20 Another aspect of the invention features Z^1 -(PML)- Z^2 containing compounds that can be used for in vivo imaging and labeling samples ex vivo, including cells; without the use of dimethylsulfoxide (DMSO) or other organic solvents (i.e. physiologic buffers or solutions) that are generally toxic to biological subjects or samples.

25 For labeling of BMs or cells, the compounds (fluorochromes) of the present invention can be incubated with BMs at various concentrations for about 5 minutes to 24 hours or more at about 4°-37°C. After the incubation the free or fluorochrome that has not been chemically linked to the BM can be removed, such as by chromatography or ultrafiltration methods that are well known in the art. For cells,
30 after the incubation, the cells can be centrifuged to create a cell pellet from which the supernatant is removed. Cells can be resuspended in culture media or physiologic saline to wash away residual, unbound or free fluorochrome. This can

be repeated several times. In this manner, cells can be labeled either by direct conjugation to internal or external cellular molecules or by non-specific cell uptake into various intracellular compartments, including but not limited to cytosol, endosomes, nucleus, golgi apparatus, and other intracellular organelles.

5 Another aspect of the invention features Z^1 -(PML)- Z^2 containing compound containing imaging probes formulated in a pharmaceutical composition suitable for administration to animal, including human, subjects. The pharmaceutical composition can include the nanoparticles and one or more stabilizers in a physiologically acceptable (relevant) carrier.

10 Another aspect of the invention features biocompatible fluorescent Z^1 -(PML)- Z^2 containing compounds formulated in pharmaceutical compositions suitable for administration to animal, including human, subjects and cells. The pharmaceutical composition can include one or more stabilizers in a physiologically acceptable (relevant) carrier.

15 Suitable examples of stabilizers for use in the methods of the present invention, include but are not limited to, low molecular weight carbohydrate, in one aspect it is a linear polyalcohol, such as sorbitol, and glycerol; or mannitol. Other low molecular weight carbohydrates, such as inositol, may also be used. Physiologically relevant carriers can include water, saline, and may further include
20 agents such as buffers, and other agents such as preservatives that are compatible for use in pharmaceutical formulations.

 The invention also features a method of gene sequence recognition using fluorescent Z^1 -(PML)- Z^2 containing compounds, labeled nucleic acid recognition molecules, including DNA, RNA, modified nucleic acid, PNA, molecular beacons,
25 aptamers; or other nucleic acid binding molecules (for example, small interfering RNA or siRNA). The method includes the use of one or more fluorescent Z^1 -(PML)- Z^2 containing compounds, together with techniques such as hybridization, ligation, cleavage, recombination, synthesis, sequencing, mutation detection, real-time polymerase chain reactions, in situ hybridization, and the use of microarrays.
30 For example, for detecting a single stranded nucleic acid (i.e., mRNA, cDNA or denatured double-stranded DNA) in a sample, via nucleic acid hybridization principles, a fluorescent Z^1 -(PML)- Z^2 containing compound chemically linked to a

single-stranded nucleic acid is contacted with a sample containing one or more single stranded nucleic acids and the fluorescence of the fluorescent Z^1 -(PML)- Z^2 containing compound is detected, wherein the presence or level of fluorescence signal emitted by the fluorescent Z^1 -(PML)- Z^2 containing compound indicates the presence or amount of nucleic acid in the sample.

The optical signal generated by the Z^1 -(PML)- Z^2 containing compounds or derivatives thereof, whether collected by tomographic, reflectance, planar, endoscopic, microscopic, surgical goggles or imager, video imaging technologies, or other methods such as microscopy including intravital and two-photon microscopy, and whether used quantitatively or qualitatively, is also considered to be an aspect of the invention.

Another aspect of the invention features a kit, which includes the Z^1 -(PML)- Z^2 containing compounds, and optionally, and instructions for using the fluorochromes or imaging probes for in vivo or in vitro imaging methods. The kit optionally may include components that aid in the use of the fluorochromes or imaging probes for the disclosed methods, such as buffers, and other formulating agents; alternatively, the kit may include medical devices that aid in the administration of the imaging probes to subjects.

The Z^1 -(PML)- Z^2 containing compounds, and pharmaceutical compositions of the present invention can be administered orally, parentally, by inhalation, topically, rectally, nasally, buccally, vaginally, or via an implanted reservoir. The term "parental administration" includes intravenous, intramuscular, subcutaneous, intraarterial, intraarticular, intrasynovial, intrasternal, intrathecal, intraperitoneal, intracisternal, intrahepatic, intralesional, intracranial and intralymphatic injection or infusion techniques. The Z^1 -(PML)- Z^2 containing compound can also be administered via catheters or through a needle to a tissue.

In one embodiment, an effective amount (which is an amount effective to cause or increase fluorescence) of the compounds of the present invention are administered. In one embodiment, between about 1 ng/kg and about 100 mg/kg, between about 100 ng/kg and 10 mg/kg, between about 1 μ g/kg and about 5 mg/kg, between about 10 μ g/kg and about 2 mg/kg, between about 50 μ g/kg and about 1 mg/kg of the compound of the present invention is administered.

Preferred Z^1 -(PML)- Z^2 containing compounds have the following properties:

- (1) high quantum yield (*e.g.*, quantum yield greater than 5% in aqueous medium),
- (2) narrow emission spectrum (*e.g.*, less than 75 nm; more preferably less than 50 nm),
- (3) spectrally separated absorption and emission spectra (*e.g.*, separated by more than 20 nm; more preferably by more than 50 nm),
- (3) have high chemical stability and photostability (*e.g.*, retain fluorescent properties after exposure to light),
- (4) are non toxic or minimally toxic to cells or subjects at doses used for imaging protocols, (as measured for example, by LD₅₀ or irritation studies, or other similar methods known in the art) and/or
- (5) have commercial viability and scalable production for large quantities (*e.g.*, gram and kilogram quantities).

The compounds of the present invention can have one or more sufficiently acidic proton that can react with a suitable organic or inorganic base to form a base addition salt. Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like, and organic bases such as alkoxides, alkyl amides, alkyl and aryl amines, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, and the like.

The compounds of the present invention having a sufficiently basic group, such as an amine can react with an organic or inorganic acid to form an acid addition salt. Acids commonly employed to form acid addition salts from compounds with basic groups are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenyl-sulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such salts include the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate,

sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, and the like.

5 The term "alkyl" as used herein means a saturated straight-chain, branched or cyclic hydrocarbon. When straight-chained or branched, an alkyl group is typically C1-C20, more typically C1-C10; when cyclic, an alkyl group is typically C3-C12, more typically C3-C7. Examples of alkyl groups include methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl and *tert*-butyl and 1,1-dimethylhexyl.

10 An "aliphatic group" is non-aromatic, and may optionally contain one or more units of unsaturation, e.g., double and/or triple bonds. An aliphatic group may be straight-chained or branched and typically contains between 1 and 12 carbon atoms, more typically between 1 and 6 carbon atoms, and even more typically between 1 and 4 carbon atoms. One or more methylene group in an aliphatic group
15 can optionally be replaced by O, S, or NH.

 As used herein the term non-aromatic carbocyclic ring or non-aromatic heterocyclic ring as used alone or as part of a larger moiety refers to a non-aromatic carbon or heteroatom containing ring which can be saturated or contain one or more units of unsaturation, having three to fourteen atoms including monocyclic and
20 polycyclic rings in which the carbocyclic or heterocyclic ring can be fused to one or more non-aromatic carbocyclic or heterocyclic rings or one or more aromatic (carbocyclic or heterocyclic) rings.

 The term "alkoxy" as used herein is represented by -OR**, wherein R** is an alkyl group as defined above.

25 The term "carbonyl" as used herein is represented by -C(=O)R**, wherein R** is an alkyl group as defined above.

 The term "aromatic group" includes carbocyclic aromatic rings and heteroaryl rings. The term "aromatic group" may be used interchangeably with the terms "aryl", "aryl ring", "aromatic ring", "aryl group" and "aromatic group".

30 Carbocyclic aromatic ring groups have only carbon ring atoms (typically six to fourteen) and include monocyclic aromatic rings such as phenyl and fused polycyclic aromatic ring systems in which a carbocyclic aromatic ring is fused to one or more

aromatic rings (carbocyclic aromatic or heteroaromatic). Examples include 1-naphthyl, 2-naphthyl, 1-anthracyl and 2-anthracyl. Also included within the scope of the term "carbocyclic aromatic ring", as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings (carbocyclic or
5 heterocyclic), such as in an indanyl, phthalimidyl, naphthimidyl, phenanthridinyl, or tetrahydronaphthyl.

The term "heteroaryl", "heteroaromatic", "heteroaryl ring", "heteroaryl group" and "heteroaromatic group", used alone or as part of a larger moiety as in "heteroaralkyl" refers to heteroaromatic ring groups having five to fourteen
10 members, including monocyclic heteroaromatic rings and polycyclic aromatic rings in which a monocyclic aromatic ring is fused to one or more other aromatic ring (carbocyclic or heterocyclic). Heteroaryl groups have one or more ring heteroatoms. Examples of heteroaryl groups include 2-furanyl, 3-furanyl, *N*-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl,
15 oxadiazolyl, oxadiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, *N*-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, *N*-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 3-pyridazinyl, 4-pyridazinyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, triazolyl, tetrazolyl, 2-thienyl, 3-thienyl, carbazolyl, benzothienyl, benzofuranyl, indolyl, quinolinyl, benzothiazole,
20 benzooxazole, benzimidazolyl, isoquinolinyl and isoindolyl. Also included within the scope of the term "heteroaryl", as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings (carbocyclic or heterocyclic).

The term non-aromatic heterocyclic group used alone or as part of a larger
25 moiety refers to non-aromatic heterocyclic ring groups having three to fourteen members, including monocyclic heterocyclic rings and polycyclic rings in which a monocyclic ring is fused to one or more other non-aromatic carbocyclic or heterocyclic ring or aromatic ring (carbocyclic or heterocyclic). Heterocyclic groups have one or more ring heteroatoms, and can be saturated or contain one or more
30 units of unsaturation. Examples of heterocyclic groups include piperidinyl, piperizinyl, pyrrolidinyl, pyrazolidinyl, imidazolidinyl, tetrahydroquinolinyl,

inodoliny, isoindoliny, tetrahydrofurany, oxazolidiny, thiazolidiny, dioxolany, dithiolany, tetrahydropyrany, dihydropyrany, azepany and azetidiny

The term "heteroatom" means nitrogen, oxygen, or sulfur and includes any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen.

5 Also the term "nitrogen" includes a substitutable nitrogen of a heteroaryl or non-aromatic heterocyclic group. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2*H*-pyrroly), NH (as in pyrrolidiny) or NR" (as in N-substituted pyrrolidiny), wherein R" is a suitable substituent for the nitrogen atom
10 in the ring of a non-aromatic nitrogen-containing heterocyclic group, as defined below. Preferably the nitrogen is unsubstituted.

A substituted aryl group as defined herein contains one or more substitutable ring atoms, such as carbon or nitrogen ring atoms. Examples of suitable substituents on a substitutable ring carbon atom of an aryl or aliphatic group include halogen
15 (e.g., -Br, Cl, I and F), -OH, C1-C4 alkyl, C1-C4 haloalkyl, -NO₂, C1-C4 alkoxy, C1-C4 haloalkoxy, -CN, -NH₂, C1-C4 alkylamino, C1-C4 dialkylamino, -C(O)NH₂, -C(O)NH(C1-C4 alkyl), -C(O)(C1-C4 alkyl), -OC(O)(C1-C4 alkyl), -OC(O)(aryl), -OC(O)(substituted aryl), -OC(O)(aralkyl), -OC(O)(substituted aralkyl), -NHC(O)H, -NHC(O)(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂, -NHC(O)O-(C1-C4 alkyl), -
20 C(O)OH, -C(O)O-(C1-C4 alkyl), -NHC(O)NH₂, -NHC(O)NH(C1-C4 alkyl), -NHC(O)N(C1-C4 alkyl)₂, -NH-C(=NH)NH₂, -SO₂NH₂ -SO₂NH(C1-C3alkyl), -SO₂N(C1-C3alkyl)₂, NHSO₂H, NHSO₂(C1-C4 alkyl) and aryl. Preferred substituents on aryl groups are as defined throughout the specification.

Examples of suitable substituents on a substitutable ring nitrogen atom of an
25 aryl group include C1-C4 alkyl, NH₂, C1-C4 alkylamino, C1-C4 dialkylamino, -C(O)NH₂, -C(O)NH(C1-C4 alkyl), -C(O)(C1-C4 alkyl), -CO₂ R**, -C(O)C(O)R**, -C(O)CH₃, -C(O)OH, -C(O)O-(C1-C4 alkyl), -SO₂NH₂ -SO₂NH(C1-C3alkyl), -SO₂N(C1-C3alkyl)₂, NHSO₂H, NHSO₂(C1-C4 alkyl), -C(=S)NH₂, -C(=S)NH(C1-C4 alkyl), -C(=S)N(C1-C4 alkyl)₂, -C(=NH)-N(H)₂, -C(=NH)-NH(C1-C4 alkyl) and
30 -C(=NH)-N(C1-C4 alkyl)₂,

Substituted alkyl, aliphatic, non-aromatic carbocyclic or heterocyclic group as defined herein contain one or more substituents. Examples of suitable

substituents for an alkyl group include those listed above for a substitutable carbon of an aryl and aliphatic and the following: =O, =S, =NNHR**, =NN(R**)2, =NNHC(O)R**, =NNHCO₂ (alkyl), =NNHSO₂ (alkyl), =NR**, spiro cycloalkyl group or fused cycloalkyl group. R** in each occurrence, independently is -H or C1-C6 alkyl. Preferred substituents on alkyl aliphatic, non-aromatic carbocyclic or heterocyclic group groups are as defined throughout the specification.

In vitro Testing and Use

The compounds (fluorochromes) and biocompatible fluorescent molecules of the present invention can be tested *in vitro* by one skilled in the art to assess its biological and performance characteristics. For instance, different types of cells grown in culture can be used to assess their biological and performance characteristics. Uptake, labeling, binding targeting or cellular localization of the fluorochromes and biocompatible fluorescent molecules can be assessed using techniques known in the art such as spectroscopy methods, fluorescent microscopy, and flow cytometry. For example, the fluorochromes and biocompatible fluorescent molecules of the present invention can be contacted with a sample for a period of time and then washed to remove any free or unbound molecules. The sample can then be viewed using a fluorescent microscope equipped with appropriate filters matched to the optical properties of the fluorochromes and biocompatible fluorescent molecules of the present invention. Fluorescent microscopy of cells in culture is also a convenient means for determining whether uptake and binding occurs in one or more subcellular compartments. Tissues, tissue sections and other types of samples such as cytospin samples can also be used in a similar manner to assess the biological and performance characteristics of the molecules. Other fluorescent detection methods including, but not limited to flow cytometry, immunoassays, hybridization assays, and microarray analysis can also be used.

Optical Imaging

The general principles of fluorescence, optical image acquisition, and image processing can be applied in the practice of the invention. For a review of optical imaging techniques, see, *e.g.*, Alfano *et al.*, *Ann. NY Acad. Sci.* 820:248-270, 1997.

An imaging system useful in the practice of this invention typically includes
5 three basic components: (1) an appropriate light source for fluorochrome and biocompatible fluorescent molecule excitation, (2) a means for separating or distinguishing emissions from light used for the excitation, and (3) a detection system to detect the optical signal emitted.

In general, the optical detection system can be viewed as including a optical
10 gathering/image forming component and a optical detection/image recording component. Although the optical detection system can be a single integrated device that incorporates both components, the optical gathering/image forming component and light detection/image recording component will be discussed separately.

A particularly useful optical gathering/image forming component is an
15 endoscope. Endoscopic devices and techniques which have been used for in vivo optical imaging of numerous tissues and organs, including peritoneum (Gahlen *et al.*, *J. Photochem. Photobiol. B* 52:131-135, 1999), ovarian cancer (Major *et al.*, *Gynecol. Oncol.* 66:122-132, 1997), colon and rectum (Mycek *et al.*, *Gastrointest. Endosc.* 48:390-394, 1998; and Stepp *et al.*, *Endoscopy* 30:379-386, 1998), bile
20 ducts (Izuishi *et al.*, *Hepatogastroenterology* 46:804-807, 1999), stomach (Abe *et al.*, *Endoscopy* 32:281-286, 2000), bladder (Kriegmair *et al.*, *Urol. Int.* 63:27-31, 1999; and Riedl *et al.*, *J. Endourol.* 13:755-759, 1999), lung (Hirsch *et al.*, *Clin Cancer Res* 7:5-220, 2001), brain (Ward, *J. Laser Appl.* 10:224-228, 1998), esophagus, and head and neck regions can be employed in the practice of the present
25 invention.

Other types of optical gathering components useful in the invention are catheter-based devices, including fiber optics devices. Such devices are particularly suitable for intravascular imaging. See, *e.g.*, Tearney *et al.*, *Science* 276:2037-2039, 1997; and *Circulation* 94:3013, 1996.

30 Still other imaging technologies, including phased array technology (Boas *et al.*, *Proc. Natl. Acad. Sci. USA* 91:4887-4891, 1994; Chance, *Ann. NY Acad. Sci.* 838:29-45, 1998), optical tomography (Cheng *et al.*, *Optics Express* 3:118-123,

1998; and Siegel *et al.*, *Optics Express* 4:287-298, 1999), intravital microscopy (Dellian *et al.*, *Br. J. Cancer* 82:1513-1518, 2000; Monsky *et al.*, *Cancer Res.* 59:4129-4135, 1999; and Fukumura *et al.*, *Cell* 94:715-725, 1998), confocal imaging (Korlach *et al.*, *Proc. Natl. Acad. Sci. USA* 96:8461-8466, 1999; 5 Rajadhyaksha *et al.*, *J. Invest. Dermatol.* 104:946-952, 1995; and Gonzalez *et al.*, *J. Med.* 30:337-356, 1999) and fluorescence molecular tomography (FMT) (Nziachristos *et al.*, *Nature Medicine* 8:757-760, 2002; United States Patent No. 6,615,063, PCT Application No. WO 03/102558, and PCT US/03/07579) can be employed in the practice of the present invention, the IVIS® Imaging System 10 (Xenogen, Alameda, CA), Maestro (CRI, Woburn, MA) the SoftScan® and the eXplore Optix™ (Advanced Research Technologies, Montreal, Canada) system can be employed in the practice of the present invention.

A suitable optical detection/image recording component, e.g., charge coupled device (CCD) systems or photographic film, can be used in the invention. The 15 choice of optical detection/image recording will depend on factors including type of optical gathering/image forming component being used. Selecting suitable components, assembling them into an optical imaging system, and operating the system is within ordinary skill in the art.

20 Diagnostic and Disease Applications and Methods

The methods of the invention can be used to determine a number of indicia, including tracking the localization of the fluorochromes and biocompatible fluorescent molecules in the subject over time or assessing changes or alterations in the metabolism and/or excretion of the molecules in the subject over time. The 25 methods can also be used to follow therapy for such diseases by imaging molecular events and biological pathways modulated by such therapy, including but not limited to determining efficacy, optimal timing, optimal dosing levels (including for individual patients or test subjects), and synergistic effects of combinations of therapy.

30 The invention can be used to help a physician or surgeon to identify and characterize areas of disease, such as arthritis, cancers and specifically colon polyps, or vulnerable plaque, to distinguish diseased and normal tissue, such as detecting

tumor margins that are difficult to detect using an ordinary operating microscope, *e.g.*, in brain surgery, help dictate a therapeutic or surgical intervention, *e.g.*, by determining whether a lesion is cancerous and should be removed or non-cancerous and left alone, or in surgically staging a disease, *e.g.*, intraoperative lymph node staging, sentinel lymph node mapping, or assessing intraoperative bleeding.

The methods of the invention can also be used in the detection, characterization and/or determination of the localization of a disease, especially early disease, the severity of a disease or a disease-associated condition, the staging of a disease, and monitoring and guiding various therapeutic interventions, such as surgical procedures, and monitoring drug therapy, including cell based therapies.

The methods of the invention can also be used in prognosis of a disease or disease condition. Examples of such disease or disease conditions include inflammation (*e.g.*, inflammation caused by arthritis, for example, rheumatoid arthritis), cancer (*e.g.*, colorectal, ovarian, lung, breast, prostate, cervical, skin, brain, gastrointestinal, mouth, esophageal, bone), cardiovascular disease (*e.g.*, atherosclerosis and inflammatory conditions of blood vessels, ischemia, stroke, thrombosis), dermatologic disease (*e.g.*, Kaposi's Sarcoma, psoriasis), ophthalmic disease (*e.g.*, macular degeneration, diabetic retinopathy), infectious disease (*e.g.*, bacterial, viral, fungal and parasitic infections, including Acquired Immunodeficiency Syndrome), immunologic disease (*e.g.*, an autoimmune disorder, lymphoma, multiple sclerosis, rheumatoid arthritis, diabetes mellitus), central nervous system disease (*e.g.*, a neurodegenerative disease, such as Parkinson's disease or Alzheimer's disease), inherited diseases, metabolic diseases, environmental diseases (*e.g.*, lead, mercury and radioactive poisoning, skin cancer), and bone-related disease (*e.g.*, osteoporosis, primary and metastatic bone tumors, osteoarthritis). The methods of the invention can therefore be used, for example, to determine the presence of tumor cells and localization of tumor cells, the presence and localization of inflammation, including the presence of activated macrophages, for instance in atherosclerosis or arthritis, the presence and localization of vascular disease including areas at risk for acute occlusion (*i.e.*, vulnerable plaques) in coronary and peripheral arteries, regions of expanding aneurysms, unstable plaque in carotid arteries, and ischemic areas. The methods and compounds (compositions) of the invention can also be used in

identification and evaluation of apoptosis, necrosis, hypoxia and angiogenesis.

Optical imaging modalities and measurement techniques include, but are not limited to, fluorescence imaging; endoscopy; fluorescence endoscopy; optical coherence tomography; transmittance imaging; time resolved transmittance imaging; 5 confocal imaging; nonlinear microscopy; photoacoustic imaging; acousto-optical imaging; spectroscopy; reflectance spectroscopy; intravital imaging; two photon imaging; interferometry; coherence interferometry; diffuse optical tomography and fluorescence molecular tomography, and measurement of light scattering, absorption, polarisation, luminescence, fluorescence lifetime, quantum yield, and 10 quenching.

The compounds (compositions) and methods of the present invention can be used in combination with other imaging compositions and methods. For example, the methods of the present invention can be used in combination with other traditional imaging modalities such as X-ray, computed tomography (CT), positron 15 emission tomography (PET), single photon computerized tomography (SPECT), and magnetic resonance imaging (MRI). For instance, the compounds (compositions) and methods of the present invention can be used in combination with CT and MR imaging to obtain both anatomical and biological information simultaneously, for example, by co-registration of a tomographic image with an image generated by 20 another imaging modality. In particular, the combination with MRI or CT is preferable, given the high spatial resolution of these imaging techniques. The compounds (compositions) and methods of the present invention can also be used in combination with X-ray, CT, PET, SPECT and MR contrast agents or the fluorescent silicon nanoparticle imaging probes of the present invention may also 25 contain components, such as iodine, gadolinium atoms and radioactive isotopes, which can be detected using CT, PET, SPECT, and MR imaging modalities in combination with optical imaging.

Kits

30 The compounds (compositions) described herein can be packaged as a kit, which may optionally include instructions for using the fluorochromes or biocompatible fluorescent molecules in various exemplary applications. Non-

limiting examples include kits that contain, *e.g.*, the compounds (compositions) in a powder or lyophilized form, and instructions for using, including reconstituting, dosage information, and storage information for *in vivo* and/or *in vitro* applications. Kits may optionally contain containers of the compounds (compositions) in a liquid
5 form ready for use, or requiring further mixing with solutions for administration. For *in vivo* applications, the kit may contain the compounds (compositions) in a dosage and form suitable for a particular application, *e.g.* a liquid in a vial, a topical creams, *etc.*

The kit can include optional components that aid in the administration of the
10 unit dose to subjects, such as vials for reconstituting powder forms, syringes for injection, customized IV delivery systems, inhalers, *etc.* The kits may be supplied in either a container which is provided with a seal which is suitable for single or multiple puncturing with a hypodermic needle (*e.g.* a crimped-on septum seal closure) while maintaining sterile integrity. Such containers may contain single or
15 multiple subject doses. Additionally, the unit dose kit can contain customized components that aid in the detection of the compounds (compositions) *in vivo* or *in vitro*, *e.g.*, specialized endoscopes, light filters. The kits may also contain instructions for preparation and administration of the compounds (compositions). The kit may be manufactured as a single use unit dose for one subject, multiple uses
20 for a particular subject; or the kit may contain multiple doses suitable for administration to multiple subjects ("bulk packaging"). The kit components may be assembled in cartons, blister packs, bottles, tubes, and the like.

Unless otherwise defined, all technical and scientific terms used herein have
25 the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated
30 by reference in their entirety. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

EXAMPLES



The following non limiting examples demonstrate the synthesis of nicotinic acid and picolinic acid derived near infrared fluorophores. Representative materials and methods that may be used in preparing the compounds of the invention are described further below. All chemicals and solvents (reagent grade) were used as supplied from the vendors cited without further purification.

The analytical and preparative HPLC methods generally utilized are:

- A Column: Agilent Zorbax 80Å, Extend C18, 4.6 x 250mm (5µm).
 Mobile Phase: Acetonitrile and 25mM triethylammonium acetate.
- B Column: Varian Dynamax, 100Å, C18, 41.4 x 250mm.
 Mobile Phase: Acetonitrile and 25mM triethylammonium acetate.
- C Column: Phenomenex Jupiter, 300Å, C18
 Mobile Phase: Acetonitrile and 25mM triethylammonium acetate.


EXAMPLE 1

Synthesis of Example 1

Part A. 6-hydrazino-1,3-naphthalene disulfonate (**I**)


6-Amino-1,3-naphthalene disulfonate disodium salt (10 g, 29 mmol) (TCI) was dissolved in 30 mL of water and added to 50 mL of water and 15 mL of concentrated hydrochloric acid. The slurry was cooled to $< 0\text{ }^{\circ}\text{C}$ in an ice/acetone bath and sodium nitrite (2.2 g, 32 mmol, Aldrich) was added in 40 mL of cold water drop-wise over 10 minutes. Stannous chloride (11 g, 58 mmol) (Aldrich) was dissolved in 30 mL water and 6 mL concentrated hydrochloric acid, cooled to $0\text{ }^{\circ}\text{C}$ and added to the reaction mixture over 10 minutes. The resulting solution was stirred and allowed to warm to room temperature over 3 hours resulting in a clear, brown solution. The solution was reduced in volume by rotary evaporation and the product precipitated by the addition of isopropanol. Product (I) was filtered, washed with isopropanol and dried in vacuum.

Part B. 2,3,3-trimethylbenzindole-5,7-disulfonate (II)




6-Hydrazino-1,3-naphthalene disulfonated (10 g, 25 mmol), isopropyl methyl ketone (12 g, 140 mmol, Aldrich) and potassium acetate (6 g, 61 mmol) were combined in 75 mL glacial acetic acid and heated to $145\text{ }^{\circ}\text{C}$ for 22 hours. The solution was cooled and the acetic acid was removed by rotary evaporation. The residue was dissolved in methanol and filtered. The product (II) was then precipitated from the methanol filtrate with isopropanol and filtered, washed with isopropanol and ether and dried in vacuum.

Part C. 2,3,3-trimethyl-1-(3-sulfonatopropyl)benzindolinium-5,7-disulfonate (III)



2,3,3-Trimethylbenzindole-6,8-disulfonate (2.2g, 5 mmol) was dissolved in 50 mL of dry DMF resulting in a clear orange solution. 1,3-propanesultone (2.8 g, 23
 5 mmol, Aldrich) was added and the solution was heated to 145 °C in a sealed tube for 15 hours, turning dark purple in color. The solution was cooled and poured into 150 mL 2-propanol. The mixture was centrifuged and the solvent decanted off. The solid product was washed on a filter with three 50 mL portions of 2-propanol followed by 50 mL of ether and dried in vacuum, resulting in 2.5 g of a dark purple
 10 solid (90%). MALDI-TOF-MS m/e 492.05 $[M]^+$ calculated for $C_{18}H_{22}NO_9S_3^+$, found 492.05.

Part D: Preparation of Compound IV



15


A 100mL round bottom flask, fitted with a reflux condenser was charged with 2,3,3-trimethyl-1-(3-sulfonatopropyl)benzindolinium-5,7-disulfonate (565 mg, 1mmol), 2-(3-hydroxycarbonyl-6-pyridyl)-malondialdehyde (98 mg, 0.5 mmol), and sodium
 20 acetate (585 mg, 7.1 mmol). Acetic anhydride (10 mL) and acetic acid (10 mL) were added to the flask, and the mixture was heated to reflux for 8 h, cooled down to room temp and 25 mL of ethyl acetate was added. The dark blue dye precipitate was collected by filtration, which was dissolved in 20 mL of water and purified by preparative reversed phase HPLC to afford the compound IV as powder. Yield, 285
 25 mg, 50%. MALDI MS: Expected, 1139.09; Found, 1139.6732

Part E: Preparation of Example 1

Compound IV (11.4 mg, 0.01 mmol) was mixed with disuccinimidyl dicarbonate (DSC, 5 mg, 0.02 mmol) and N,N-dimethylaminopyridine (DMAP, 2 mg, 0.016 mmol) in 500 μ L of dry DMF, and heated to 70 $^{\circ}$ C. After 30 min cold ethyl acetate (500 μ L) was added to the reaction mixture to precipitate the product out, which was filtered, dried under vacuum and stored at 4 $^{\circ}$ C.


10 **EXAMPLE 2**

Synthesis of Example 2



15

Part A: Preparation of Compound V



20 Compound V was prepared from potassium, 2,3,3-trimethyl-1-ethyl-benzindolinium-5,7-disulfonate (184 mg, 0.423 mmol), 2-(3-hydroxycarbonyl-6-pyridyl)-malondialdehyde (40 mg, 0.21 mmol), and sodium acetate (300 mg, 3.66

mmol) mixed in 6 mL of acetic anhydride and 6 mL of acetic acid. After heating at 125 °C for 16 hours, the product was precipitated from ethyl acetate and purified by reversed phase HPLC. The product in a powder form was obtained after speed vacuum drying of the HPLC fractions. MALDI. m/z calculated (found) for M^+ :


5 952.05 (952.16). Yield, 140 mg, 70%.

Part B: Preparation of Example 2

One equivalent of Compound V was mixed with two equivalents of disuccinimidyl dicarbonate (DSC) and one equivalent of N,N-dimethylaminopyridine (DMAP) in 10 0.5 mL of DMF, and heated to 70 °C. The reaction was complete within 30 min; the desired product was precipitated from the DMF solution by the addition of ethyl acetate, filtered, dried under vacuum and stored at 4 °C.


15 EXAMPLE 3

Synthesis of Example 3



20

Part A: Preparation of Compound VI



Compound VI was prepared from 2,3,3-Trimethyl-1-(3-sulfonatopropyl)-3H-benzindolinium-5,7-disulfonate (101 mg, 0.18 mmol), 2-(2-hydroxycarbonyl-6-pyridyl)-malonaldialdehyde (17.4 mg, 0.09 mmol), and sodium acetate (88 mg, 1.07 mmol) mixed in 6 mL of acetic anhydride and 6 mL of acetic acid. After heating at 125 °C for 4 hours, the product was precipitated from ethyl acetate and purified by reversed phase HPLC. The product was obtained in a powder form after speed vacuum drying of the HPLC fractions. MALDI. *m/z* calculated (found) for (M+1)⁺: 1140.33 (1140.13). Yield, 67 mg, 65%.

10

Part B: Preparation of Example 3


One equivalent of compound VI was mixed with two equivalents of disuccinimidyl dicarbonate (DSC) and one equivalent of N,N-dimethylaminopyridine (DMAP) in 0.5 mL of DMF, and heated to 70 °C. The progress of the reaction was monitored by HPLC. The reaction was complete within 30 min, following which the desired product was precipitated from the DMF solution by the addition of ethyl acetate, filtered, dried under vacuum and stored at 4 °C.

15

EXAMPLE 4

20

Synthesis of Example 4



Example 4 is prepared from 2,3,3-Trimethyl-1-(ethyl)-3H-benzindolinium-5,7-disulfonate (0.18 mmol), 2-(2-hydroxycarbonyl-6-pyridyl)-malonaldialdehyde (0.09 mmol), and sodium acetate (88 mg, 1.07 mmol) mixed in 6 mL of acetic anhydride and 6 mL of acetic acid. After heating at 125 °C for 4 hours, the product is


25

precipitated from ethyl acetate, purified by reversed phase HPLC and obtained in a powder form after speed vacuum drying of the HPLC fractions.

EXAMPLE 5

5

Synthesis of Example 5




- 10 2,3,3-Trimethyl-1-(3-sulfonatopropyl)-indolinium-5-sulfonate (115 mg, 0.28 mmol), 2-(3-hydroxycarbonyl-6-pyridyl)-malonaldialdehyde (27 mg, 0.14 mmol), and sodium acetate (80 mg, 1.0 mmol) were mixed in 5 mL of acetic anhydride and 5 mL of acetic acid. After heating at 125 °C for 2 hours, reaction mixture was cooled down and the product was precipitated from ethyl acetate, followed by purification by reversed phase HPLC. The product was obtained in a powder form after speed vacuum drying of the HPLC fractions. Yield, 87 mg, 70%. Abs. max (water): 635 nm; Em. Max (water): 653 nm.

EXAMPLE 6

20

Synthesis of Example 6




- 25 2-Methyl-1-(3-sulfonatopropyl)-benzothiazolium inner salt (150 mg, 0.55 mmol) and 2-(3-hydroxycarbonyl-6-pyridyl)-malonaldialdehyde (52 mg, 0.27 mmol) were added to a mixture of 2-methoxy ethanol and toluene (2:1, v/v) and heated to reflux

at 150 °C attached to a Dean-Stark condenser. After 4 hours, heating was stopped, cooled down and solvents were removed by rotary evaporation. The concentrated blue dye was purified using preparative HPLC. The product was obtained in a powder form after drying of the HPLC fractions on speed vacuum. Yield, 95 mg, 50%. Abs. max (water): 643 nm; Em. Max (water): 663 nm.

EXAMPLE 7

Synthesis of Example 7




10

10 mg of the NHSE of Compound IV was dissolved in 100uL dry DMF, to which was added a solution of 3-azidopropylamine (5 mg) in 20 uL dry DMSO, and the mixture was rotated at room temperature for one hour. 1 mL ether was added to the reaction mixture, and centrifuged for 10 minutes. The supernatant solution was discarded, and the residue was dried on speed vac for 5 minutes, redissolved in water and purified on a RPC18 semi-prep column. The fraction corresponding to the product (Example 7) was collected, and dried on speedvac which yielded 7.2 mg (60%). It was characterized by MALDI. Calculated: 1222.34. Found: 1222.54

20

EXAMPLE 8

Synthesis of Example 8




25

10 mg of the NHSE of Compound IV was dissolved in 100uL dry DMF, to which was added a solution of 3-propargylamine (5 mg) in 10 uL dry DMSO, and the mixture was rotated at room temperature for one hour. 1 mL ether was added to the reaction mixture, and centrifuged for 10 minutes. The supernatant solution was discarded, and the residue was dried on speed vac for 5 minutes, redissolved in water and purified on a RPC18 semi-prep column. The fraction corresponding to the product (Example 8) was collected, and dried on speedvac. Yield: 6 mg (52%). It was characterized by MALDI. Calculated: 1177.32. Found: 1177.21

EXAMPLE 9

Synthesis of Example 9



17 mg of the NHSE of Compound IV was dissolved in 250uL dry DMF, to which was added a solution of 2-(2-aminoethyl-dithio)pyridine hydrochloride (11 mg) in 10 uL dry DMSO and 4 uL of triethylamine, and the mixture was rotated at room temperature for an overnight. 1 mL ethyl acetate was added to the reaction mixture, and centrifuged for 10 minutes. The supernatant solution was discarded, and the residue was dried on speed vac for 5 minutes, redissolved in water and purified on a RPC18 semi-prep column. The fraction corresponding to the product (Example 9) was collected, and dried on speedvac. Yield: 10 mg (65%). It was characterized by MALDI. Calculated: 1308.52. Found: 1308.54

EXAMPLE 10**Cell Labeling**

Mouse splenocytes were prepared as a single cell suspension, and the T cell subpopulation within the splenocyte preparation were enriched by passage over a column that removes B cells and macrophages (R& D kit, Mouse T-cell enrichment columns, MTCC500). T cells were centrifuged to generate a cell pellet of 10^7 cells. The supernatant was removed from the cell pellet, and a solution of 10 mg/ml of compound of Example 1 in 100 uL was added. Cells were incubated at room temperature for 5 minutes, followed by 2 rounds of centrifugation and resuspension in physiologic buffer to wash away unbound Example 1. Cells were assessed by fluorescence microscopy.

EXAMPLE 11**Cell Labeling and In Vivo Imaging**

Mouse 4T1 breast adenocarcinoma cells were centrifuged to generate a cell pellet of 10^7 cells. The supernatant was removed from the cell pellet, and a solution of 10 mg/ml Example 1 in 100 uL was added. Cells were incubated at room temperature for 5 minutes, followed by 2 rounds of centrifugation and resuspension in physiologic buffer to wash away unbound Example 1. Cells were assessed by fluorescence microscopy. Cells were injected intravenously into mice at 5×10^5 cells per mouse, and live mice were imaged by fluorescent molecular tomography immediately after injection and 24 hours after injection. As 4T1 cells primarily metastasize to the lungs, lung fluorescence is quantified.

EXAMPLE 12

A solution of Example 1 was chemically linked to an amine presenting surface of nanoparticles to yield a biocompatible fluorescent molecule for in vivo optical imaging. The tumor cell line HT-29 (human colon carcinoma/HTB-38) was obtained from ATCC (Manassas, VA). HT-29 cells were grown in McCoy's supplemented with 10% FBS at 37 °C in a humidified atmosphere containing 5% CO₂. Exponentially growing cells were trypsinized and re-suspended in Hank's Balanced Salt Solution at a concentration of 3×10^7 cells/ml. Female NU/NU mice 6-


8 weeks old (Charles River Laboratory, Wilmington, MA) were injected subcutaneously with 3×10^6 HT-29 cells bilaterally in the first mammary fat pads. One week later, when tumors were approximately 30 mm^3 , mice were injected intravenously with the fluorescent molecule, (in $150 \mu\text{l}$ of $1 \times \text{PBS}$) and imaged after
5 24 hrs on a fluorescence reflectance system (FRI, Kodak 2000MM) system. Results are shown in FIG. 1

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled
10 in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

CLAIMS

What is claimed is:

1. A compound of the formula (2):



(2)

or a salt thereof, wherein:

X is independently selected from the group consisting of C(CH₂Y₁)(CH₂Y₂), O, S, and Se;

Y₁ and Y₂ are independently selected from the group consisting of H, C₁-C₂₀ aliphatic group and a C₁-C₂₀ aliphatic group substituted with -OR*, N(R*)₂ or -SR*;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;


R₁ is selected from the group consisting of H, (CH₂)_xCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R₄ is selected from the group consisting of H, (CH₂)_xCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R₂ and R₃ are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group.

2. The method of Claim 1, wherein the wherein:
- a) the carboxyl substituent on Q is selected from the group consisting of an ester, an activated ester; or
- 5 b) the carboxyl substituent on Q is selected from the group consisting of CO-Obenzotriazolyl, CO-ON-succinimidyl, CO-OTetrafluorophenyl, CO-Opentafluorophenyl, CO-Oimidazolyl, and CO-Op-nitrophenyl; and
- c) the carbonyl substituent on Q is in the form of a carbonyl halide.
- 10 3. The compound of Claim 2, wherein Q is a carboxyl substituted nitrogen containing heterocyclic ring.
4. The compound of Claim 3, wherein Q is selected from the group consisting of carboxyl substituted pyridine, pyrimidone, pyrazine, and pyridazine.
- 15 5. The compound of Claim 4, wherein Q is carboxyl substituted pyridine.
6. The compound of Claim 5, wherein Q is selected from a group consisting of isonicotinic acid, nicotinic acid and picolinic acid.
- 20 7. The compound of Claim 6, wherein Q is represented by a structural formula selected from a group consisting of:



25

8. The compound of Claim 1, wherein Q is a carbonyl substituted nitrogen containing 6-membered heterocyclic ring
- 30 9. The compound of Claim 8, wherein Q is a carbonyl substituted pyridine.


10. The compound of Claim 1, wherein at least one of the moieties R_1 to R_3 is, or contains a sulphonic acid moiety or a sulphonate moiety.

5


11. The compound of Claim 1, wherein R_1 and R_4 are independently $-H$, $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$.

12. The compound of Claim 1, having any of the formulae:

10




14. The compound of Claim 1, having any of the formulae:



15

15. A compound of the formula (2):



(2)

or a salt thereof, wherein:

X is independently selected from the group consisting of
 5 C(CH₂Y₁)(CH₂Y₂), O, S, and Se;

Y₁ and Y₂ are independently selected from the group consisting of H,
 C₁-C₂₀ aliphatic group and a C₁-C₂₀ aliphatic group substituted with -OR*,
 N(R*)₂ or -SR*;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-
 10 condensed ring;

R₁ is selected from the group consisting of (CH₂)_xCH₃, (CH₂)_nSO₃⁻
 and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an
 integer selected from 2 to 6;

R₄ is selected from the group consisting of (CH₂)_xCH₃, (CH₂)_nSO₃⁻
 15 and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an
 integer selected from 2 to 6;

R₂ and R₃ are independently selected from the group consisting of H,
 carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide,
 hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted
 20 with a carboxyl group or 6-membered heteroaryl ring substituted with a
 carbonyl group.

16. The method of Claim 15, wherein the wherein:

- 25 a) the carboxyl substituent on Q is selected from the group consisting of
 an ester, an activated ester; or
 b) the carboxyl substituent on Q is selected from the group consisting of
 CO-Obenzotriazolyl, CO-ON-succinimidyl, CO-Otetrafluorophenyl,

CO-Opentafluorophenyl, CO-Oimidazolyl, and CO-Op-nitrophenyl;
and

c) the carbonyl substituent on Q is in the form of a carbonyl halide.

5 17. The compound of Claim 15, wherein Q is a carboxyl substituted nitrogen containing heterocyclic ring.

18. The compound of Claim 17, wherein Q is selected from the group consisting of carboxyl substituted pyridine, pyrimidone, pyrazine, and pyridazine.


10

19. The compound of Claim 18, wherein Q is carboxyl substituted pyridine.

20. The compound of Claim 19, wherein Q is selected from a group consisting of isonicotinic acid, nicotinic acid and picolinic acid.

15

21. The compound of Claim 20, wherein Q is represented by a structural formula selected from a group consisting of:



20

22. The compound of Claim 15, wherein Q is a carbonyl substituted nitrogen containing 6-membered heterocyclic ring

23. The compound of Claim 22, wherein Q is a carbonyl substituted pyridine.


25

24. The compound of Claim 15, wherein at least one of the moieties R_1 to R_3 is, or contains a sulphonic acid moiety or a sulphonate moiety.

25. The compound of Claim 15, wherein R_1 and R_4 are independently $(CH_2)_nSO_3^-$ or $(CH_2)_nSO_3H$.


30

26. The compound of Claim 15 having any of the formulae:




5

27. The compound of Claim 15 having any of the formulae:



28. A biocompatible fluorescent molecule comprising one or more biomolecules
10 chemically linked to a compound represented by formula (2):



(2)

or a salt thereof, wherein:

- 15 X is independently selected from the group consisting of
C(CH₂Y₁)(CH₂Y₂), O, S, and Se;

Y_1 and Y_2 are independently selected from the group consisting of H, C_1 - C_{20} aliphatic group and a C_1 - C_{20} aliphatic group substituted with $-OR^*$, $N(R^*)_2$ or $-SR^*$;

5 W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;


R_1 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

10 R_4 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_2 and R_3 are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxy, a sulphonic acid moiety and a sulphonate moiety;

15 Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group.

29. A biocompatible fluorescent molecule comprising a biomolecule chemically
20 linked to a compound represented by formula (2):



(2)

or a salt thereof, wherein:

25 X is independently selected from the group consisting of $C(CH_2Y_1)(CH_2Y_2)$, O, S, and Se;

Y_1 and Y_2 are independently selected from the group consisting of H, C_1 - C_{20} aliphatic group and a C_1 - C_{20} aliphatic group substituted with $-OR^*$, $N(R^*)_2$ or $-SR^*$;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;

R_1 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_4 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;


R_2 and R_3 are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group;

wherein the compound has an absorption and emission maxima between about 500 nm and about 900 nm

30. The compound of Claim 29, wherein the fluorochrome has an absorption and emission maxima between about 600 nm and 800 nm.

31. A biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound represented by formula (2):



(2)

or a salt thereof, wherein:

X is independently selected from the group consisting of $C(CH_2Y_1)(CH_2Y_2)$, O, S, and Se;

Y_1 and Y_2 are independently selected from the group consisting of H, C_1 - C_{20} aliphatic group and a C_1 - C_{20} aliphatic group substituted with $-OR^*$, $N(R^*)_2$ or $-SR^*$;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;


R_1 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_4 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_2 and R_3 are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group; wherein the biocompatible fluorescent molecule is activated after target interaction.

32. A biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound represented by formula (2):



(2)

or a salt thereof, wherein:

X is independently selected from the group consisting of $C(CH_2Y_1)(CH_2Y_2)$, O, S, and Se;

Y_1 and Y_2 are independently selected from the group consisting of H, C_1 - C_{20} aliphatic group and a C_1 - C_{20} aliphatic group substituted with $-OR^*$, $N(R^*)_2$ or $-SR^*$;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;

R_1 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R_4 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$ and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;


R_2 and R_3 are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group;

wherein the biocompatible fluorescent molecule has a high binding affinity to a target.

33. A method of in vivo optical imaging, the method comprising:

- (a) administering to a subject a biocompatible fluorescent molecule comprising one or more biomolecules chemically linked to a compound represented by formula (2):



(2)

or a salt thereof, wherein:

X is independently selected from the group consisting of
 $C(CH_2Y_1)(CH_2Y_2)$, O, S, and Se;

5 Y_1 and Y_2 are independently selected from the group consisting of H,
C₁-C₂₀ aliphatic group and a C₁-C₂₀ aliphatic group substituted with -OR*,
N(R*)₂ or -SR*;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-
condensed ring;

10 R_1 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$
and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an
integer selected from 2 to 6;

R_4 is selected from the group consisting of H, $(CH_2)_xCH_3$, $(CH_2)_nSO_3^-$
and $(CH_2)_nSO_3H$, wherein x is an integer selected from 0 to 6 and n is an
integer selected from 2 to 6;

15 R_2 and R_3 are independently selected from the group consisting of H,
carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide,
hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

20 Q is selected from a group consisting of a heteroaryl ring substituted
with a carboxyl group or 6-membered heteroaryl ring substituted with a
carbonyl group


- (b) allowing time for the biocompatible fluorescent molecule to
distribute within the subject or to contact or interact with a biological
target;
- (c) illuminating the subject with light of a wavelength absorbable by the
25 biocompatible fluorescent molecule; and
- (d) detecting the optical signal emitted by the biocompatible fluorescent
molecule.

34. The method of Claim 33, wherein the signal emitted by the biocompatible
30 fluorescent molecule is used to construct an image.

35. The method of Claim 34, wherein steps (a) - (d) are repeated at predetermined intervals thereby allowing for evaluation of the emitted signals of the biocompatible fluorescent molecule in the subject over time.
- 5 36. The method of Claim 35, wherein the subject is an animal or human.
37. The method of Claim 36, wherein in step (a) two or more fluorescent molecule whose signal properties are distinguishable are administered to a subject.
- 10 38. The method of Claim 37, wherein the illuminating and detecting steps are performed using an endoscope, catheter, tomographic system, hand-held optical imaging system, surgical goggles, or intra-operative microscope.
- 15 39. The method of Claim 38, wherein the presence, absence, or level of signal emitted by the fluorescent molecule is indicative of a disease state.
40. The method of Claim 39, wherein the method is used to detect and monitor disease.
- 20 41. The method of Claim 40, wherein the disease is selected from the group consisting of cancer, cardiovascular diseases, neurodegenerative diseases, immunologic diseases, autoimmune diseases, respiratory diseases, metabolic diseases, inherited diseases, infectious diseases, bone diseases, and environmental diseases.
- 25 42. The method of Claim 41, wherein prior to step (a) the fluorescent molecule of claim 1 is mixed with cells to label the cells and the resulting labeled cells are administered to the subject in step (a).
- 30

43. The method of Claim 42, wherein the signal emitted by the fluorescent molecule is used for monitoring trafficking and localization of cells or evaluation a cell therapy.

5 44. A biocompatible fluorescent molecule comprising a biomolecule chemically linked to a compound represented by formula (2):



(2)

10 or a salt thereof, wherein:

X is independently selected from the group consisting of C(CH₂Y₁)(CH₂Y₂), O, S, and Se;

Y₁ and Y₂ are independently selected from the group consisting of H, C₁-C₂₀ aliphatic group and a C₁-C₂₀ aliphatic group substituted with -OR*, N(R*)₂ or -SR*;

W represents a benzo-condensed, a naphtho-condensed or a pyrido-condensed ring;

R₁ is selected from the group consisting of H, (CH₂)_xCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R₄ is selected from the group consisting of H, (CH₂)_xCH₃, (CH₂)_nSO₃⁻ and (CH₂)_nSO₃H, wherein x is an integer selected from 0 to 6 and n is an integer selected from 2 to 6;

R₂ and R₃ are independently selected from the group consisting of H, carboxylate, carboxylic acid, carboxylic ester, amine, amide, sulfonamide, hydroxyl, alkoxyl, a sulphonic acid moiety and a sulphonate moiety;

Q is selected from a group consisting of a heteroaryl ring substituted with a carboxyl group or 6-membered heteroaryl ring substituted with a carbonyl group;

wherein the biomolecule is a labeled cell.

1/1



FIG. 1